

http://www.nlm.nih.gov/medlineplus/news/fullstory_74357.html

Most Visited Getting Started Latest Headlines

Skip navigation

MedlinePlus
Trusted Health Information for You

A service of the U.S. NATIONAL LIBRARY OF MEDICINE
and the NATIONAL INSTITUTES OF HEALTH

Search MedlinePlus

About MedlinePlus | Site Map | FAQs | Contact Us

Home Health Topics Drugs & Supplements Encyclopedia Dictionary **News** Directories Other Resources

HealthDay

Page Title

HS Sports Shoulder Injuries More Likely in Boys
Player-to-player contact in competition behind almost 60% of cases, study finds

Printer-friendly version E-mail to a friend

Author
HealthDay
By Robert Preidt

Date
Monday, January 26, 2009

URL

Organization

MONDAY, Jan. 26 (HealthDay News) -- Male high school athletes in the United States are more likely than girls to suffer shoulder injuries.

The researchers looked at football, soccer, volleyball, basketball, wrestling, baseball and softball.

Overall, shoulder injuries accounted for just 8 percent of all injuries suffered by high school athletes. However, shoulder injuries were fairly common in predominately male sports such as baseball (18 percent of all injuries), wrestling (18 percent), and football (12 percent). Boys had higher shoulder injury rates than girls, particularly in soccer and baseball/softball.

Shoulder injuries were three times more likely to occur during competition than in practice.

Player-to-player contact was associated with almost 60 percent of high-school athletes' shoulder injuries. The most common shoulder injuries included sprains and strains (37 percent), dislocations and separations (24 percent), contusions (12 percent), and fractures (7 percent).

Done

Six percent of shoulder injuries required surgery, and dislocations and separations accounted for more than half of all shoulder surgeries.

"Wrestling shoulder injuries were most likely to require surgery, with almost one in 10 requiring such procedures," study co-author Dawn Comstock, principal investigator in the Center for Injury Research and Policy at Nationwide Children's Hospital in Columbus, Ohio, said in a hospital news release.

"Even more importantly, in all sports, almost one in four athletes missed at least three weeks of the season following a shoulder injury. This underscores the importance of preventing shoulder injuries before they occur," said Comstock, a faculty member at the Ohio State University College of Medicine.

The study, published online in the *Journal of Athletic Training*, involved an analysis of data from the 2005-2007 National High School Sports Injury Surveillance System.

HealthDay

Copyright (c) 2009 ScoutNews, LLC. All rights reserved.

[More News on this Date](#)

Related MedlinePlus Pages:

- [Shoulder Injuries and Disorders](#)
- [Sports Injuries](#)
- [Teen Health](#)

Home | Health Topics | Drugs & Supplements | **Organization** | News | Directories | Other Resources

Date

Disclaimers | Copyright | Privacy | Accessibility | Quality Guidelines
U.S. National Library of Medicine, 8600 Rockville Pike, Bethesda, MD 20894
National Institutes of Health | Department of Health & Human Services

Date last updated: 27 January 2009